

The robust noncontextuality inequalities in the simplest scenario

Matthew F. Pusey

Perimeter Institute for Theoretical Physics, 31 Caroline Street North, Waterloo, ON N2L 2Y5, Canada*

(Dated: June 12, 2015)

Contextuality is the leading notion of non-classicality for a single system. However, experimental demonstrations of contextuality have generally been limited by conceptual difficulties. Here I focus on the simplest non-trivial case, four preparations and two tomographically complete binary measurements. Exploiting a subtle connection to the CHSH scenario gives eight non-linear inequalities which are together necessary and sufficient for the experimental statistics to admit a preparation noncontextual model. No fixed “operational equivalences” are required. Violating such an inequality is therefore arguably the most direct route to an experimental refutation of noncontextuality.

The gold standard for an experiment that defies classical explanation is the violation of a Bell inequality [1, 2]. In the case of a single quantum system, this is not a possibility, and so attention has focussed on contextuality.

Contextuality was first identified by Bell, Kochen and Specker [3, 4]. Whilst this was a profound insight into quantum mechanics, the definition they used is not amenable to principled experimental test (see [5, 6] and references therein). The definition was later refined and generalised by Spekkens [7]. As shown in [8], this definition, or even just one component of it known as *preparation noncontextuality*, is more suited to experiment.

However, [8] used an assumption that two preparation procedures were indistinguishable, which was not satisfied exactly in the reported experiment and never will be in any experiment. Here I show how this problem can be eliminated by providing a full characterisation of the preparation noncontextual statistics in the simplest scenario to which the concept applies. The shift in approach is that, with the help of tomographically complete measurements, indistinguishable preparation procedures are inferred from the statistics, rather than posited *a priori*.

Definitions. Consider an experiment where one implements a preparation procedure $\mathcal{P}_i$ followed by a measurement procedure $\mathcal{M}_j$ with outcome k , characterised by the probabilities $P(k|\mathcal{P}_i, \mathcal{M}_j)$. An *ontological model* seeks to explain these results via an *ontic state* λ that screens off the preparation from the measurement result:

$$P(k|\mathcal{P}_i, \mathcal{M}_j) = \int P(k|\lambda, \mathcal{M}_j) \mu_i(\lambda) d\lambda, \quad (1)$$

where we use the shorthand $\mu_i(\lambda) = P(\lambda|\mathcal{P}_i)$.

The explanation proffered by an ontological model is compelling only if it does justice to important features of the observed statistics. For example, in a bipartite scenario, Bell’s locally causal models would provide a natural explanation for the observed no-signalling [9].

Preparation noncontextuality concerns a closely related feature, namely *operational equivalence* among mixtures of preparations. Suppose there exists probability distributions $\{p_i\}$ and $\{q_i\}$ such that for all j, k

$$\sum_i p_i P(k|\mathcal{P}_i, \mathcal{M}_j) = \sum_i q_i P(k|\mathcal{P}_i, \mathcal{M}_j). \quad (2)$$

Then we say that the probabilistic mixtures, which might be written $\sum_i p_i \mathcal{P}_i$ and $\sum_i q_i \mathcal{P}_i$, are operationally equivalent. In principle it is possible that somebody invents a new measurement procedure $\mathcal{M}'$ such that $\sum_i p_i P(k|\mathcal{P}_i, \mathcal{M}') \neq \sum_i q_i P(k|\mathcal{P}_i, \mathcal{M}')$, in which case the apparent operational equivalence would evaporate. For the time being we will assume no such $\mathcal{M}'$ exists. In the language of quantum theory, where preparations $\mathcal{P}_i$ are associated with density operators ρ_i , we are assuming that $\sum_i p_i \rho_i = \sum_i q_i \rho_i$. This follows from eq. (2) if and only if the POVMs associated with the $\mathcal{M}_j$ are *tomographically complete* [10] or *fiducial* [11] for the density operators spanned by the ρ_i . This crucial issue will be returned to later.

What can explain the inability of any measurement to distinguish $\sum_i p_i \mathcal{P}_i$ from $\sum_i q_i \mathcal{P}_i$? The most natural explanation is that this “distinction without a difference” is no distinction at all:

$$\sum_i p_i \mu_i(\lambda) = \sum_i q_i \mu_i(\lambda). \quad (3)$$

The inference from the operational equivalence (2) to the ontic equivalence (3) constitutes the assumption of preparation noncontextuality [7]. (For comparison, measurement noncontextuality is the assumption that measurements that cannot be distinguished by the statistics for any preparation are equivalent in the ontological model.)

The simplest scenario. Suppose we consider only a single measurement $\mathcal{M}_1$. Then the simple “ $\lambda = k$ ” model $P(k|\lambda, \mathcal{M}_1) = \delta_{k\lambda}$, $\mu_i(\lambda) = P(k = \lambda|\mathcal{P}_i, \mathcal{M}_1)$ is manifestly preparation noncontextual. Hence we need to consider at least two measurements, the simplest case being two binary (two-outcome) measurements.

In this scenario the operational probabilities for a single preparation $\mathcal{P}_i$ are given by the 2-dimensional real vector $\vec{\mathcal{P}}_i = (P(0|\mathcal{P}_i, \mathcal{M}_0) - P(1|\mathcal{P}_i, \mathcal{M}_0), P(0|\mathcal{P}_i, \mathcal{M}_1) - P(1|\mathcal{P}_i, \mathcal{M}_1))$ which (along with normalisation) fixes all 4 probabilities. Hence eq. (2) has the geometric form $\sum_i p_i \vec{\mathcal{P}}_i = \sum_i q_i \vec{\mathcal{P}}_i$. The convex hull of any three $\vec{\mathcal{P}}_i$ is a simplex: a point, a line segment, or (the generic case) a triangle. Consider the ontological model in which there is an ontic state for each extreme point of the simplex, with $\mu_i(\lambda) = 1$ for the one or more $\vec{\mathcal{P}}_i$ located at the extreme




FIG. 1. Example statistics. The preparations have been labelled in accordance with the conventions that $\vec{P}_0$ is opposite $\vec{P}_3$ and $\{\vec{P}_0 - \vec{P}_3, \vec{P}_2 - \vec{P}_1\}$ is positively oriented. Also shown is $\vec{c}$ as defined by eq. (6).

point $\vec{\lambda}$, and $p(k|\lambda, \mathcal{M}_i)$ defined by the coordinates of the extreme point in the obvious way. Expanding eq. (2) as

$$\sum_{i,\lambda} p_i \mu_i(\lambda) \vec{\lambda} = \sum_{i,\lambda} q_i \mu_i(\lambda) \vec{\lambda} \quad (4)$$

the fact that every point in a simplex has a unique convex decomposition into extremal points immediately gives eq. (3), so our model is preparation noncontextual.

Hence the simplest scenario in which one can demonstrate noncontextuality, and the focus of most of this article, is four preparations and two binary measurements.

Labelling conventions. Denoting the four preparations $\{\mathcal{P}_0, \mathcal{P}_1, \mathcal{P}_2, \mathcal{P}_3\}$, the $\vec{P}_i = (x_i, y_i)$ must be the vertices of a quadrilateral, since any degeneracy will lead to a simplex and hence an immediate preparation noncontextual model by the above argument. As in fig. 1, we adopt the conventions that $\vec{P}_0$ is opposite to $\vec{P}_3$, and the $\vec{P}_0 - \vec{P}_3$ and $\vec{P}_2 - \vec{P}_1$ diagonals are positively oriented:

$$\begin{vmatrix} x_0 - x_3 & x_2 - x_1 \\ y_0 - y_3 & y_2 - y_1 \end{vmatrix} > 0. \quad (5)$$

A pivotal equivalence. One example of an operational equivalence is given by the point $\vec{c}$ at which the $\{\vec{P}_0, \mathcal{P}_3\}$ diagonal intersects $\{\vec{P}_1, \mathcal{P}_2\}$ diagonal, giving probabilities p, q such that

$$p\vec{P}_0 + (1-p)\vec{P}_3 = q\vec{P}_1 + (1-q)\vec{P}_2 = \vec{c}. \quad (6)$$

Preparation noncontextuality then demands that

$$p\mu_0(\lambda) + (1-p)\mu_3(\lambda) = q\mu_1(\lambda) + (1-q)\mu_2(\lambda). \quad (7)$$

I will now show that in the current scenario, this single equivalence is in fact sufficient for a model to be preparation noncontextual.

Suppose we have a preparation contextual model, i.e. there exists p_i, q_i such that eq. (2) holds yet eq. (3)

fails. We want to prove that eq. (7) must also fail. The first step is to show that eq. (3) must fail for some p'_i, q'_i with $\sum_i p'_i \vec{P}_i = \sum_i q'_i \vec{P}_i = \vec{c}$. To see this, denote $\vec{p} = \sum_i p_i \vec{P}_i = \sum_i q_i \vec{P}_i$, and notice that since $\vec{c}$ is in the interior of the quadrilateral $\vec{c} = \sum_i r_i \vec{P}_i + r_* \vec{p}$ for some probability distribution $\{r_0, r_1, r_2, r_3, r_*\}$ with $r_* > 0$. But then $p'_i = r_i + r_* p_i$ and $q'_i = r_i + r_* q_i$ give the required instance, with the failure of (3) ensured by the fact it fails for the p_i, q_i and $r_* > 0$.

Now I will argue that there exist probabilities s, t such that $\{p'_0, p'_1, p'_2, p'_3\} = \{sp, (1-s)q, (1-s)(1-q), s(1-p)\}$ and $\{q'_0, q'_1, q'_2, q'_3\} = \{tp, (1-t)q, (1-t)(1-q), t(1-p)\}$, i.e. that $\sum_i p'_i \vec{P}_i$ amounts to preparing $p\mathcal{P}_0 + (1-p)\mathcal{P}_2$ with probability s and $q\mathcal{P}_1 + (1-q)\mathcal{P}_2$ with probability $(1-s)$ (similarly for the q'_i with t in place of s). If that is the case then (7) implies $\sum_i p'_i \mu_i(\lambda) = \sum_i q'_i \mu_i(\lambda)$ and so the failure of the latter requires the failure of the former, which is what we wanted to prove.

To find s and t it is useful to make an affine transformation to a new co-ordinate system in which $\vec{c} = (0, 0)$, $\vec{P}_0 = (1-p, 0)$ and $\vec{P}_1 = (0, 1-q)$. Then $p\vec{P}_0 + (1-p)\vec{P}_3 = \vec{c}$ gives $\vec{P}_3 = (-p, 0)$ and $q\vec{P}_1 + (1-q)\vec{P}_2 = \vec{c}$ gives $\vec{P}_2 = (0, -q)$. Now $\sum_i p'_i \vec{P}_i = \vec{c}$ becomes $(p'_0(1-p) - p'_3p, p'_1(1-q) - p'_2q) = (0, 0)$. Defining $s = p'_0/p \geq 0$ we have $p'_3 = s(1-p)$. Similarly defining $\bar{s} = p'_1/q \geq 0$ we have $p'_2 = \bar{s}(1-q)$. $\sum_i p_i = 1$ gives $s + \bar{s} = 1$, and repeating the same argument with the q'_i gives t .

The connection to CHSH. It was first shown by Barrett that the existence of a preparation noncontextual model for a single system implies the existence of a locally causal model for any bipartite scenario involving that system [12], in other words, any bipartite proof of Bell's theorem is a proof of preparation contextuality. The converse is not expected to hold in general (although certain proofs of preparation contextuality can be converted into bipartite proofs of Bell's theorem [13]). Nevertheless, with the reduction of the previous section in hand, Barrett's argument can be extended to see that the existence of a preparation noncontextual model for four preparations and two binary measurements is *equivalent* to the existence of a Bell local model in the scenario considered by CHSH [14].

In the relevant Bell scenario two parties choose between two measurements, their choices labelled a and b . They both obtain a binary outcome, labelled x and y . Their statistics $P(a, b|x, y)$ are related to the preparation noncontextuality scenario by

$$\begin{pmatrix} P(0, k|0, j) & P(1, k|0, j) \\ P(0, k|1, j) & P(1, k|1, j) \end{pmatrix} = \begin{pmatrix} pP(k|\mathcal{P}_0, \mathcal{M}_j) & (1-p)P(k|\mathcal{P}_3, \mathcal{M}_j) \\ qP(k|\mathcal{P}_1, \mathcal{M}_j) & (1-q)P(k|\mathcal{P}_2, \mathcal{M}_j) \end{pmatrix}. \quad (8)$$

These statistics are normalised, and are no-signalling due to the operational equivalence (6). If we have

a preparation noncontextual model, then set $\mu(\lambda) = p\mu_0(\lambda) + (1-p)\mu_3(\lambda)$,

$$\begin{pmatrix} P_A(0|\lambda, 0) & P_A(1|\lambda, 0) \\ P_A(0|\lambda, 1) & P_A(1|\lambda, 1) \end{pmatrix} = \begin{pmatrix} p\mu_0(\lambda)/\mu(\lambda) & (1-p)\mu_3(\lambda)/\mu(\lambda) \\ q\mu_1(\lambda)/\mu(\lambda) & (1-q)\mu_2(\lambda)/\mu(\lambda) \end{pmatrix} \quad (9)$$

(which is normalised by eq. (2)), and $P_B(k|\lambda, j) = P(k|\lambda, \mathcal{M}_j)$. Then eq. (1) gives the locally causal model

$$p(a, b|x, y) = \int P_A(a|\lambda, x)P_B(b|\lambda, y)\mu(\lambda)d\lambda. \quad (10)$$

If, conversely, we start with a locally causal model, then inverting eq. (9) gives an ontological model for $P(k|\mathcal{P}_i, \mathcal{M}_j)$, where eq. (7) is guaranteed by the normalisation of the locally causal model. Since we have seen that eq. (7) is sufficient for a preparation noncontextual model, we have established the desired equivalence.

Fine [15] has shown that the eight versions of the CHSH inequality [14] are necessary and sufficient for the existence of a locally causal model. Hence given $P(k|\mathcal{P}_i, \mathcal{M}_j)$, one can calculate the corresponding Bell scenario probabilities (8) and then use the eight CHSH inequalities to determine whether or not a preparation noncontextual model exists.

A closed expression. This above argument completely characterises the noncontextual statistics in our scenario. However, it might appear that if one were to calculate p and q explicitly and substitute (8) into the CHSH inequalities the result would be extremely convoluted. In fact it can be written in a remarkably simple form, thanks to following lemma.

Lemma 1. *Suppose $\vec{P}_i = (x_i, y_i)$ for $i = 0, 1, 2, 3$ satisfy eq. (5), and p and q are defined as the solutions of eq. (6). Then for any real numbers $\{z_0, z_1, z_2, z_3\}$,*

$$pz_0 + (1-p)z_3 \leq qz_1 + (1-q)z_2 \quad (11)$$

if and only if

$$\begin{vmatrix} x_0 & y_0 & z_0 & 1 \\ x_1 & y_1 & z_1 & 1 \\ x_2 & y_2 & z_2 & 1 \\ x_3 & y_3 & z_3 & 1 \end{vmatrix} \leq 0. \quad (12)$$

Furthermore, equality in (11) and (12) is also equivalent.

Geometrically this lemma concerns a tetrahedron with vertices (x_i, y_i, z_i) . Equation (11) asks whether, when the 0-3 edge meets the 1-2 edge in the (x, y) -plane, it is below in the z -direction. Subject to the convention (5) that is equivalent to the statement (12) about the signed volume of the tetrahedron. A purely algebraic proof can be given as follows.

Proof. Denoting the LHS of (5) by D , Cramer's rule gives

$$p = \begin{vmatrix} x_2 - x_3 & x_2 - x_1 \\ y_2 - y_3 & y_2 - y_1 \end{vmatrix} / D, \quad (13)$$

$$q = \begin{vmatrix} x_0 - x_3 & x_2 - x_3 \\ y_0 - y_3 & y_2 - y_3 \end{vmatrix} / D. \quad (14)$$

Substituting these into (11) and multiplying through by $D > 0$ gives, upon expanding all the determinants, (12). The reader can avoid an algebraic quagmire by referring to the Mathematica notebook provided on the arXiv. $\square$

If we substitute eq. (8) into the CHSH inequalities we obtain (11) with $z_0 = c_0x_0 + d_0y_0 - 1$, $z_1 = -c_1x_1 - d_1y_1 + 1$, $z_2 = c_1x_2 + d_1y_2 + 1$, $z_3 = -c_0x_3 - d_0y_3 - 1$, where (c_0, d_0, c_1, d_1) is a column of

$$\begin{pmatrix} 1 & 1 & 1 & 1 & -1 & -1 & -1 & -1 \\ 1 & 1 & -1 & -1 & 1 & 1 & -1 & -1 \\ 1 & -1 & 1 & -1 & 1 & -1 & 1 & -1 \\ -1 & 1 & 1 & -1 & 1 & -1 & -1 & 1 \end{pmatrix}. \quad (15)$$

(There is one column for each version of the CHSH inequality.) Substituting the z_i into (12) then gives the desired inequality. For example the $\langle A_0B_0 \rangle + \langle A_0B_1 \rangle + \langle A_1B_0 \rangle - \langle A_1B_1 \rangle \leq 2$ version of the CHSH inequality corresponds to the first column of (15), and (12) becomes

$$\begin{vmatrix} x_0 & y_0 & x_0 + y_0 - 1 & 1 \\ x_1 & y_1 & -x_1 + y_1 + 1 & 1 \\ x_2 & y_2 & x_2 - y_2 + 1 & 1 \\ x_3 & y_3 & -x_3 - y_3 - 1 & 1 \end{vmatrix} \leq 0. \quad (16)$$

(Subject to (5), this can still serve as a Bell inequality, but now in terms of $p(a|b, x, y)$ rather than $p(a, b|x, y)$.)

Quantum violation. Suppose $\mathcal{M}_0$ and $\mathcal{M}_1$ correspond to X and Z measurements of a qubit. Let $\mathcal{P}_0$ correspond to preparing the $+1$ eigenstate of $(X + Z)/\sqrt{2}$ and $\mathcal{P}_3$ the -1 eigenstate. Similarly let $\{\mathcal{P}_1, \mathcal{P}_2\}$ be the $\{+1, -1\}$ eigenstates of $(X - Z)/\sqrt{2}$. Denoting $v = 1/\sqrt{2}$ the LHS of (16) is

$$\begin{vmatrix} v & v & 2v - 1 & 1 \\ v & -v & 1 - 2v & 1 \\ -v & v & 1 - 2v & 1 \\ -v & -v & 2v - 1 & 1 \end{vmatrix} = 16v^2(2v - 1) \approx 3.31. \quad (17)$$

This is the same proof of the preparation contextuality of a qubit that appeared in [8]. However, that proof assumed $\frac{1}{2}\mathcal{P}_1 + \frac{1}{2}\mathcal{P}_3 = \frac{1}{2}\mathcal{P}_1 + \frac{1}{2}\mathcal{P}_2$, which will never hold exactly in a realistic experiment. Since no such assumption entered into (16), the proof presented here is more experimentally robust.

Tomographic completeness. A new difficulty arises in the present approach, namely our assumption that $\mathcal{M}_0$ and $\mathcal{M}_1$ are tomographically complete for the $\mathcal{P}_i$. An obvious objection in the qubit example of the previous section is the Y measurement. Suppose we have




FIG. 2. Dealing with failure of tomographic completeness. The statistics for three binary measurements $\{\mathcal{M}_0, \mathcal{M}_1, \mathcal{M}'\}$ are defined by a 3-dimensional vector for each preparation. Two views of this space are shown. The four non-planar preparations in brown, together with additional preparation in blue, imply the existence of the four planar preparations in green. The statistics for the green preparations can be calculated with simple trigonometry and then tested against the inequalities derived here for two binary measurements.

a preparation noncontextual model for three measurements $\{\mathcal{M}_0, \mathcal{M}_1, \mathcal{M}'\}$, and consider a set of preparations that all give the same probability for $\mathcal{M}'$. Since eq. (2) holds for all three measurements if and only if it holds for $\{\mathcal{M}_0, \mathcal{M}_1\}$, the problem reduces to finding a preparation noncontextual model for $\mathcal{M}_0$ and $\mathcal{M}_1$. Since the quantum states mentioned above all give uniformly random outcomes for the Y measurement, there is in principle nothing wrong with using (16).

In practice, however, the preparations in a real experiment will each give slightly different probabilities for the Y measurement. The simplest way to deal with this is to add a fifth preparation that gives one outcome of the Y measurement with high probability (e.g. the $+1$ eigenstate of Y). If, as in fig. 2, we then consider a plane perpendicular to the Y axis with the four original preparations on one side and the new preparations on the other, convexity implies the existence of, and determines the probabilities for, four preparations in the plane that can then be tested against (16). Notice there is no need to actually implement the four new preparations. Any further measurements that reveal small amounts of information about the preparations can be dealt with similarly.

This idea has been generalised [16] into a technique for identifying experimental violations of other noncontextuality inequalities, even those requiring fixed equivalences.

Conclusions. The eight inequalities derived here fully classify the preparation noncontextual statistics in the simplest non-trivial scenario. No assumptions of determinism or unattainable operational equivalences were made. No new assumptions on the representation of approximately operationally equivalent procedures [17, 18] were made either. Hence a violation can only be explained by a failure of noncontextuality or a failure of the tomographic completeness of the measurements. Since

observed failures of the latter can be dealt with using convexity, non-classicality may be left as the only plausible explanation.

The main direction for future theoretical work is to classify scenarios with more preparations and measurements. Failures of tomographic completeness could then be dealt more elegantly than above, by fully incorporating the extra procedures into the contextuality scenario. It would also be interesting to apply similar ideas to measurement and transformation noncontextuality.

Thanks to Rob Spekkens for invaluable discussions, and to Jon Barrett for sharing his result [12] and allowing me to include it here. Research at Perimeter Institute is supported in part by the Government of Canada through NSERC and by the Province of Ontario through MRI.

* m@physics.org

- [1] J. S. Bell, *Physics* **1**, 195 (1964).
- [2] N. Brunner, D. Cavalcanti, S. Pironio, V. Scarani, and S. Wehner, *Rev. Mod. Phys.* **86**, 419 (2014), arXiv:1303.2849.
- [3] J. S. Bell, *Rev. Mod. Phys.* **38**, 447 (1966).
- [4] S. Kochen and E. Specker, *Indiana Univ. Math. J.* **17**, 59 (1968).
- [5] J. Barrett and A. Kent, *Stud. Hist. Phil. Mod. Phys.* **35**, 151 (2004), arXiv:quant-ph/0309017.
- [6] R. W. Spekkens, *Found. Phys.* **44**, 1125 (2014), arXiv:1312.3667.
- [7] R. W. Spekkens, *Phys. Rev. A* **71**, 052108 (2005), arXiv:quant-ph/0406166.
- [8] R. W. Spekkens, D. H. Buzacott, A. J. Keehn, B. Toner, and G. J. Pryde, *Phys. Rev. Lett.* **102**, 010401 (2009), arXiv:0805.1463.
- [9] C. J. Wood and R. W. Spekkens, *New J. Phys.* **17**, 033002 (2015), arXiv:1208.4119.
- [10] V. Bužek, R. Derka, G. Adam, and P. Knight, *Ann. Phys.* **266**, 454 (1998).
- [11] L. Hardy, “Quantum theory from five reasonable axioms,” (2001), arXiv:quant-ph/0101012.
- [12] J. Barrett, “Any bipartite Bell-type proof of non-locality can be converted into a proof preparation contextuality,” (2004), unpublished note received in 2010.
- [13] M. S. Leifer and O. J. E. Maroney, *Phys. Rev. Lett.* **110**, 120401 (2013), arXiv:1208.5132.
- [14] J. F. Clauser, M. A. Horne, A. Shimony, and R. A. Holt, *Phys. Rev. Lett.* **23**, 880 (1969).
- [15] A. Fine, *Phys. Rev. Lett.* **48**, 291 (1982).
- [16] M. D. Mazurek, M. F. Pusey, R. Kunjwal, K. J. Resch, and R. W. Spekkens, “An experimental test of noncontextuality without unwarranted idealizations,” (2015), arXiv:1505.06244.
- [17] A. Winter, *J. Phys. A: Math. Theor.* **47**, 424031 (2014), arXiv:1408.0945.
- [18] J. V. Kujala, E. N. Dzhafarov, and J.-Å. Larsson, “Necessary and sufficient conditions for contextuality in a broad class of quantum mechanical systems,” (2014), arXiv:1412.4724.